Charles Moore, M.D. L. Henry Pham, M.D. Phyllis C. Chang, M.D.

Christopher J. Derbes, M.D. Joshua Sultz, M.D. David Case, M.D.

William H. Lovett, III, PA-C Kayce Kieschnick, PA-C Hannah Helm, PA-C

# **PATIENT INSTRUCTIONS – UPPER EGD:**

#### 2 WEEKS BEFORE PROCEDURE

Discontinue all DIET PILLS and APPETITE SUPPRESSANTS (Phentermine or Phendimetrazine) –
please let your scheduler know if you are currently taking these types of medication.
This is for your safety under anesthesia.

### **5 DAYS** BEFORE PROCEDURE

Discontinue all Oral Iron Supplements. Discontinue use of Wegovy, Ozempic, and other glp1 agonists.

### **3 DAYS** BEFORE PROCEDURE

- o If you are taking Blood Thinners, contact the prescribing provider for instructions.
- Discontinue all Herbal supplements, Vitamins, Anti-inflammatories (Advil, Motrin, Ibuprofen, Aleve).
   It is okay to take Tylenol.
- o Avoid all Beans, Peas, Corn, Popcorn and Grapes.

### AFTER MIDNIGHT

# NO SMOKING, CHEWING TOBACCO, or ANY TOBACCO PRODUCTS.

NO gum, mints, candy or cough drops.

### **DAY OF PROCEDURE**

You May take your approved prescribed morning medications with small sips of water. (These include blood pressure meds, anti-seizure meds, & anti-rejection meds). Finish these at least 1 hour prior to starting the prep.

Exact arrival times will be given the day before your procedure. Arrival Time is 30-45 minutes before your procedure time and is subject to change within a 4 hour window. Your driver should arrive with you.

### **IMPORTANT TIPS:**

- 1. <u>You MUST have someone to drive you home.</u> Your driver should be prepared to remain in the waiting area or in their vehicle during your procedure.
- 2. You may apply Vaseline or Diaper Rash Ointment to the rectal area if you experience discomfort from frequent stools.
- 3. IF YOU GET NAUSEOUS, or have had a weight loss procedure, you may start the Prep Earlier so that you can drink slower and take frequent breaks. You must drink the entire prep as instructed, regardless of your weight or what your stools look like.
- 4. If you have problems tolerating the prep, or you are not getting Results, please call (979) 776-4600 and ask to speak to a triage nurse (or after hour physician).

# **COMMON CANCELLATION REASONS:**

- No Driver
- Drinking fluids within 4 hours of procedure time.
- Smoking or using tobacco products the day of procedure.
- Taking Diet Meds within 2 weeks of procedure

Charles Moore, M.D. L. Henry Pham, M.D. Phyllis C. Chang, M.D.

Christopher J. Derbes, M.D. Joshua Sultz, M.D. David Case, M.D.

William H. Lovett, III, PA-C Kayce Kieschnick, PA-C Hannah Helm, PA-C

# **MEDICAL CONDITIONS:**

If you are affected by any of the conditions or medications listed below, please follow these Instructions:

	-
DIABETES	<b>ORAL MEDICATIONS:</b> On the day PRIOR to your procedure take your morning
	doses as usual. Do NOT take your evening dose the day prior to your procedure. Do
	NOT take your diabetic medications on the morning of your procedure.
	<b>INSULIN:</b> On the day PRIOR to your procedure take your morning dose of insulin
	as usual. On the evening prior to your procedure take only half of the regular evening
	dose of insulin. <b>Do NOT take your insulin the morning of your procedure.</b>
	Check your blood sugar frequently while taking the prep solution and the morning of
	your procedure. If you develop symptoms associated with an abnormally low blood
	sugar please contact the doctor at (979) 776-4600.
ASPIRIN	If you are taking aspirin PRESCRIBED by your MD, please continue to take it. If
	you are taking aspirin on your own, please stop taking aspirin 5 days before your
	procedure.
COUMADIN,	Ask the physician who prescribed your medicine how to take it before and after your
PLAVIX, ELEQUIS,	procedure. If you cannot contact your physician, call us several days before your
XARELTO OR	procedure.
OTHER BLOOD	
THINNERS	

Please refer to your pre-procedural instruction page or contact us for further information or clarification.

If you do not hear from the Facility your procedure is scheduled at, please reach out to them the day before your procedure.

PROCEDURES CANCELLED LESS THAN 24 HOURS IN ADVANCE WILL BE SUBJECT TO A \$100 CANCELLATION FEE.